

Potentially $K_m - G$ -graphical Sequences: A Survey *

Chunhui Lai, Lili Hu

Department of Mathematics and Information Science,

Zhangzhou Teachers College,

Zhangzhou, Fujian 363000, P. R. of CHINA.

zjlaichu@public.zzptt.fj.cn (Chunhui Lai, Corresponding author)

jackey2591924@163.com (Lili Hu)

Abstract

The set of all non-increasing nonnegative integers sequence $\pi = (d(v_1), d(v_2), \dots, d(v_n))$ is denoted by NS_n . A sequence $\pi \in NS_n$ is said to be graphic if it is the degree sequence of a simple graph G on n vertices, and such a graph G is called a realization of π . The set of all graphic sequences in NS_n is denoted by GS_n . A graphical sequence π is potentially H -graphical if there is a realization of π containing H as a subgraph, while π is forcibly H -graphical if every realization of π contains H as a subgraph. Let K_k denote a complete graph on k vertices. Let $K_m - H$ be the graph obtained from K_m by removing the edges set $E(H)$ of the graph H (H is a subgraph of K_m). This paper summarizes briefly some recent results on potentially $K_m - G$ -graphic sequences and give a useful classification for determining $\sigma(H, n)$.

Key words: graph; degree sequence; potentially $K_m - G$ -graphic sequences

AMS Subject Classifications: 05C07, 05C35

1 Introduction

The set of all non-increasing nonnegative integer sequence $\pi = (d(v_1), d(v_2), \dots, d(v_n))$ is denoted by NS_n . A sequence $\pi \in NS_n$ is said to be graphic

*Project Supported by NSF of Fujian(Z0511034), Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering", Project of Fujian Education Department and Project of Zhangzhou Teachers College.

if it is the degree sequence of a simple graph G on n vertices, and such a graph G is called a realization of π . The set of all graphic sequences in NS_n is denoted by GS_n . A graphical sequence π is potentially H -graphical if there is a realization of π containing H as a subgraph, while π is forcibly H -graphical if every realization of π contains H as a subgraph. If π has a realization in which the $r+1$ vertices of largest degree induce a clique, then π is said to be potentially A_{r+1} -graphic. Let $\sigma(\pi) = d(v_1) + d(v_2) + \dots + d(v_n)$, and $[x]$ denote the largest integer less than or equal to x . We denote $G + H$ as the graph with $V(G + H) = V(G) \cup V(H)$ and $E(G + H) = E(G) \cup E(H) \cup \{xy : x \in V(G), y \in V(H)\}$. Let K_k , C_k , T_k , and P_k denote a complete graph on k vertices, a cycle on k vertices, a tree on $k+1$ vertices, and a path on $k+1$ vertices, respectively. Let F_k denote the friendship graph on $2k+1$ vertices, that is, the graph of k triangles intersecting in a single vertex. For $0 \leq r \leq t$, denote the generalized friendship graph on $kt - kr + r$ vertices by $F_{t,r,k}$, where $F_{t,r,k}$ is the graph of k copies of K_t meeting in a common r set. We use the symbol Z_4 to denote $K_4 - P_2$. Let $K_m - H$ be the graph obtained from K_m by removing the edges set $E(H)$ of the graph H (H is a subgraph of K_m).

Given a graph H , what is the maximum number of edges of a graph with n vertices not containing H as a subgraph? This number is denoted $ex(n, H)$, and is known as the Turán number. In terms of graphic sequences, the number $2ex(n, H) + 2$ is the minimum even integer l such that every n -term graphical sequence π with $\sigma(\pi) \geq l$ is forcibly H -graphical. Erdős, Jacobson and Lehel [13] first consider the following variant: determine the minimum even integer l such that every n -term graphical sequence π with $\sigma(\pi) \geq l$ is potentially H -graphical. We denote this minimum l by $\sigma(H, n)$. Erdős, Jacobson and Lehel [13] showed that $\sigma(K_k, n) \geq (k-2)(2n-k+1) + 2$ and conjectured that equality holds. They proved that if π does not contain zero terms, this conjecture is true for $k=3$, $n \geq 6$. The conjecture is confirmed in [19] and [43-46]. Li et al. [46] and Mubayi [55] also independently determined the values $\sigma(K_r, 2k)$ for any $k \geq 3$ and $n \geq k$. Li and Yin [51] further determined $\sigma(K_r, n)$ for $r \geq 7$ and $n \geq 2r+1$. The problem of determining $\sigma(K_r, n)$ is completely solved.

Gould, Jacobson and Lehel [19] also proved that $\sigma(pK_2, n) = (p-1)(2n-p) + 2$ for $p \geq 2$; $\sigma(C_4, n) = 2\lceil \frac{3n-1}{2} \rceil$ for $n \geq 4$. Lai [29] gave a lower bound of $\sigma(C_k, n)$ and proved that $\sigma(C_5, n) = 4n-4$ for $n \geq 5$ and $\sigma(C_6, n) = 4n-2$ for $n \geq 7$. Lai [32] proved that $\sigma(C_{2m+1}, n) = m(2n-m-1) + 2$, for $m \geq 2, n \geq 3m$; $\sigma(C_{2m+2}, n) = m(2n-m-1) + 4$, for $m \geq 2, n \geq 5m-2$. Li and Luo [41] gave a lower bound of $\sigma({}_3C_l, n)$ and determined $\sigma({}_3C_l, n)$, $4 \leq l \leq 6, n \geq l$. Li, Luo and Liu [42] determined $\sigma({}_3C_l, n)$ for $3 \leq l \leq 8$ and $n \geq l$. and $\sigma({}_3C_9, n)$ for $n \geq 12$. Li and Yin [48] determined $\sigma({}_3C_l, n)$ for n sufficiently large. Yin, Li and Chen [68] determined $\sigma({}_kC_l, n)$, $l \geq 7, 3 \leq k \leq l$. Chen and Yin [9] determined the values

$\sigma(W_5, n)$ for $n \geq 11$ where W_r is a wheel graph on r vertices. For $r \times s$ complete bipartite graph $K_{r,s}$, Gould, Jacobson and Lehel[19] determined $\sigma(K_{2,2}, n)$. Yin et al. [63,65,69,70] determined $\sigma(K_{r,s}, n)$ for $s \geq r \geq 2$ and sufficiently large n . For $r \times s \times t$ complete 3-partite graph $K_{r,s,t}$, Erdős, Jacobson and Lehel[13] determined $\sigma(K_{1,1,1}, n)$. Lai[30] determined $\sigma(K_{1,1,2}, n)$. Yin[58] and Lai[34] independently determined $\sigma(K_{1,1,3}, n)$. Chen[7] determined $\sigma(K_{1,1,t}, n)$ for $t \geq 3$, $n \geq 2\lceil \frac{(t+5)^2}{4} \rceil + 3$. Chen[5] determined $\sigma(K_{1,2,2}, n)$ for $5 \leq n \leq 8$ and $\sigma(K_{2,2,2}, n)$ for $n \geq 6$. Let K_s^t denote the complete t partite graph such that each partite set has exactly s vertices. Guantao Chen, Michael Ferrara, Ronald J. Gould, John R. Schmitt[11] showed that $\sigma(K_s^t, n) = \pi(K_{(t-2)s} + K_{s,s}, n)$ and obtained the exact value of $\sigma(K_j + K_{s,s}, n)$ for n sufficiently large. Consequently, they obtained the exact value of $\sigma(K_s^t, n)$ for n sufficiently large. For $n \geq 5$, Ferrara, Jacobson and Schmitt[17] determined $\sigma(F_k, n)$ where F_k denotes the graph of k triangles intersecting at exactly one common vertex. In[16], Ferrara, Gould and Schmitt determined a lower bound for $\sigma(K_s^t, n)$ where K_s^t denotes the complete multipartite graph with t partite sets each of size s , and proved equality in the case $s = 2$. They also provided a graph theoretic proof of the value of $\sigma(K_t, n)$. Michael J. Ferrara[15] determined $\sigma(H, n)$ for the graph $H = K_{m_1} \cup K_{m_2} \cup \dots \cup K_{m_k}$ where n is sufficiently large integer. Ferrara, M., Jacobson, M., Schmitt, J. and Siggers M.[18] determined $\sigma(K_{s,t}, m, n)$, $\sigma(P_t, m, n)$ and $\sigma(C_{2t}, m, n)$ where $\sigma(H, m, n)$ is the minimum integer k such that every bigraphic pair $S = (A, B)$ with $|A| = m$, $|B| = n$ and $\sigma(S) \geq k$ is potentially H -bigraphic. For an arbitrarily chosen H , Schmitt, J.R. and Ferrara, M.[56] gave a good lower bound of $\sigma(H, n)$. Yin and Li[67] determined $\sigma(K_{r_1, r_2, \dots, r_l, r, s}, n)$ for sufficiently large n . Moreover, Yin, Chen and Schmitt[62] determined $\sigma(F_{t,r,k}, n)$ for $k \geq 2$, $t \geq 3$, $1 \leq r \leq t-2$ and sufficiently large, where $F_{t,r,k}$ denotes the graph of k copies of K_t meeting in a common r set. Gupta, Joshi and Tripathi [20] gave a necessary and sufficient condition for the existence of a tree of order n with a given degree set. Yin [59] gave a new necessary and sufficient condition for π to be potentially K_{r+1} -graphic. Jiong-sheng Li and Jianhua Yin [50] gave a survey on graphical sequences.

2 Potentially $K_m - G$ -graphical Sequences

Let H be a graph with m vertices, then $H = K_m - (K_m - H)$. Let $G = K_m - H$, then $\sigma(H, n) = \sigma(K_m - G, n)$. If the Problem 1 - 5 in the Open Problems section be solved, then the problem of determining $\sigma(H, n)$ is completely solved. We think the Problem 3, 4 is a useful classification for determining $\sigma(K_m - G, n)$.

Gould, Jacobson and Lehel[19] pointed out that it would be nice to see

where in the range for $3n - 2$ to $4n - 4$, the value $\sigma(K_4 - e, n)$ lies. Later, Lai[30] proved that

Theorem 1. For $n = 4, 5$ and $n \geq 7$

$$\sigma(K_4 - e, n) = \begin{cases} 3n - 1 & \text{if } n \text{ is odd} \\ 3n - 2 & \text{if } n \text{ is even.} \end{cases}$$

For $n = 6$, S is a 6-term graphical sequence with $\sigma(S) \geq 16$, then either there is a realization of S containing $K_4 - e$ or $S = (3^6)$. (Thus $\sigma(K_4 - e, 6) = 20$).

Huang[26] gave a lower bound of $\sigma(K_m - e, n)$. Yin, Li and Mao[71] and Huang[27] independently determined the values $\sigma(K_5 - e, n)$ as following.

Theorem 2. If $n \geq 5$, then

$$\sigma(K_5 - e, n) = \begin{cases} 5n - 7, & \text{if } n \text{ is odd} \\ 5n - 6, & \text{if } n \text{ is even} \end{cases}$$

Lai[35-36] determined $\sigma(K_5 - C_4, n)$, $\sigma(K_5 - P_3, n)$ and $\sigma(K_5 - P_4, n)$.

Theorem 3. For $n \geq 5$, $\sigma(K_5 - C_4, n) = \sigma(K_5 - P_3, n) = \sigma(K_5 - P_4, n) = 4n - 4$.

Yin and Li[66] gave a good method of determining the values $\sigma(K_{r+1} - e, n)$ (In fact, Yin and Li[66] also determining the values $\sigma(K_{r+1} - ke, n)$ for $r \geq 2$ and $n \geq 3r^2 - r - 1$).

Theorem 4. Let $n \geq r + 1$ and $\pi = (d_1, d_2, \dots, d_n) \in GS_n$ with $d_{r+1} \geq r$. If $d_i \geq 2r - i$ for $i = 1, 2, \dots, r - 1$, then π is potentially A_{r+1} -graphic.

Theorem 5. Let $n \geq 2r + 2$ and $\pi = (d_1, d_2, \dots, d_n) \in GS_n$ with $d_{r+1} \geq r$. If $d_{2r+2} \geq r - 1$, then π is potentially A_{r+1} -graphic.

Theorem 6. Let $n \geq r + 1$ and $\pi = (d_1, d_2, \dots, d_n) \in GS_n$ with $d_{r+1} \geq r - 1$. If $d_i \geq 2r - i$ for $i = 1, 2, \dots, r - 1$, then π is potentially $K_{r+1} - e$ -graphic.

Theorem 7. Let $n \geq 2r + 2$ and $\pi = (d_1, d_2, \dots, d_n) \in GS_n$ with $d_{r-1} \geq r$. If $d_{2r+2} \geq r - 1$, then π is potentially $K_{r+1} - e$ -graphic.

Theorem 8. If $r \geq 2$ and $n \geq 3r^2 - r - 1$, then

$$\sigma(K_{r+1} - ke, n) = \begin{cases} (r - 1)(2n - r) - (n - r) + 1, & \text{if } n - r \text{ is odd} \\ (r - 1)(2n - r) - (n - r) + 2, & \text{if } n - r \text{ is even} \end{cases}$$

After reading[66], Yin[72] determined the values $\sigma(K_{r+1} - K_3, n)$ for $r \geq 3, n \geq 3r + 5$.

Theorem 9. If $r \geq 3$ and $n \geq 3r + 5$, then $\sigma(K_{r+1} - K_3, n) = (r - 1)(2n - r) - 2(n - r) + 2$.

Determining $\sigma(K_{r+1} - H, n)$, where H is a tree on 4 vertices is more useful than a cycle on 4 vertices (for example, $C_4 \not\subset C_i$, but $P_3 \subset C_i$ for $i \geq 5$). So, after reading [66] and [72], Lai and Hu [38] determined $\sigma(K_{r+1} - H, n)$ for $n \geq 4r + 10, r \geq 3, r + 1 \geq k \geq 4$ and H be a graph on k vertices which containing a tree on 4 vertices but not containing a cycle on 3 vertices and $\sigma(K_{r+1} - P_2, n)$ for $n \geq 4r + 8, r \geq 3$.

Theorem 10. If $r \geq 3$ and $n \geq 4r + 8$, then $\sigma(K_{r+1} - P_2, n) = (r - 1)(2n - r) - 2(n - r) + 2$.

Theorem 11. If $r \geq 3, r + 1 \geq k \geq 4$ and $n \geq 4r + 10$, then $\sigma(K_{r+1} - H, n) = (r - 1)(2n - r) - 2(n - r)$, where H is a graph on k vertices which contains a tree on 4 vertices but not contains a cycle on 3 vertices.

There are a number of graphs on k vertices which containing a tree on 4 vertices but not containing a cycle on 3 vertices (for example, the cycle on k vertices, the tree on k vertices, and the complete 2-partite graph on k vertices, etc).

Lai and Sun [39] determined $\sigma(K_{r+1} - (kP_2 \cup tK_2), n)$ for $n \geq 4r + 10, r + 1 \geq 3k + 2t, k + t \geq 2, k \geq 1, t \geq 0$.

Theorem 12. If $n \geq 4r + 10, r + 1 \geq 3k + 2t, k + t \geq 2, k \geq 1, t \geq 0$, then $\sigma(K_{r+1} - (kP_2 \cup tK_2), n) = (r - 1)(2n - r) - 2(n - r)$.

To now, the problem of determining $\sigma(K_{r+1} - H, n)$ for H not containing a cycle on 3 vertices and sufficiently large n has been solved.

Lai [37] determined $\sigma(K_{r+1} - Z, n)$ for $n \geq 5r + 19, r + 1 \geq k \geq 5, j \geq 5$ and Z is a graph on k vertices and j edges which contains a graph Z_4 but not contain a cycle on 4 vertices. In the same paper, the author also determined the values of $\sigma(K_{r+1} - Z_4, n)$, $\sigma(K_{r+1} - (K_4 - e), n)$, $\sigma(K_{r+1} - K_4, n)$ for $n \geq 5r + 16, r \geq 4$.

Theorem 13. If $r \geq 4$ and $n \geq 5r + 16$, then

$$\begin{aligned} \sigma(K_{r+1} - K_4, n) &= \sigma(K_{r+1} - (K_4 - e), n) = \\ \sigma(K_{r+1} - Z_4, n) &= \begin{cases} (r - 1)(2n - r) - 3(n - r) + 1, & \text{if } n - r \text{ is odd} \\ (r - 1)(2n - r) - 3(n - r) + 2, & \text{if } n - r \text{ is even} \end{cases} \end{aligned}$$

Theorem 14. If $n \geq 5r + 19, r + 1 \geq k \geq 5$, and $j \geq 5$, then

$$\sigma(K_{r+1} - Z, n) = \begin{cases} (r - 1)(2n - r) - 3(n - r) - 1, & \text{if } n - r \text{ is odd} \\ (r - 1)(2n - r) - 3(n - r) - 2, & \text{if } n - r \text{ is even} \end{cases}$$

where Z is a graph on k vertices and j edges which contains a graph Z_4 but not contains a cycle on 4 vertices.

There are a number of graphs on k vertices and j edges which contains a graph Z_4 but not contains a cycle on 4 vertices. (for example, the graph obtained by $C_3, C_{i_1}, C_{i_2}, \dots, C_{i_p}$ intersecting in a single vertex ($i_j \neq 4, j = 1, 2, 3, \dots, p$) (if $i_j = 3, j = 1, 2, 3, \dots, p$, then this graph is the friendship graph F_{p+1}), the graph obtained by $C_3, P_{i_1}, P_{i_2}, \dots, P_{i_p}$ intersecting in a single vertex, ($i_1 \geq 1$), the graph obtained by $C_3, P_{i_1}, C_{i_2}, \dots, C_{i_p}$ ($i_j \neq 4, j = 2, 3, \dots, p, i_1 \geq 1$) intersecting in a single vertex, etc)

Lai and Yan[40] proved that

Theorem 15. If $n \geq 5r + 18, r + 1 \geq k \geq 7$, and $j \geq 6$, then

$$\sigma(K_{r+1} - U, n) = \begin{cases} (r-1)(2n-r) - 3(n-r) - 1, \\ \text{if } n-r \text{ is odd} \\ (r-1)(2n-r) - 3(n-r), \\ \text{if } n-r \text{ is even} \end{cases}$$

where U is a graph on k vertices and j edges which contains a graph $(K_3 \cup P_3)$ but not contains a cycle on 4 vertices and not contains Z_4 .

There are a number of graphs on k vertices and j edges which contains a graph $(K_3 \cup P_3)$ but not contains a cycle on 4 vertices and not contains Z_4 . (for example, $C_3 \cup C_{i_1} \cup C_{i_2} \cup \dots \cup C_{i_p}$ ($i_j \neq 4, j = 2, 3, \dots, p, i_1 \geq 5$), $C_3 \cup P_{i_1} \cup P_{i_2} \cup \dots \cup P_{i_p}$ ($i_1 \geq 3$), $C_3 \cup P_{i_1} \cup C_{i_2} \cup \dots \cup C_{i_p}$ ($i_j \neq 4, j = 2, 3, \dots, p, i_1 \geq 3$), etc)

A harder question is to characterize the potentially H -graphic sequences without zero terms. Luo [53] characterized the potentially C_k -graphic sequences for each $k = 3, 4, 5$.

Theorem 16. Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 3$. Then π is potentially C_3 -graphic if and only if $d_3 \geq 2$ except for 2 case: $\pi = (2^4)$ and $\pi = (2^5)$.

Theorem 17. Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 4$. Then π is potentially C_4 -graphic if and only if the following conditions hold:

- (1) $d_4 \geq 2$. (2) $d_1 = n - 1$ implies $d_2 \geq 3$
- (3) If $n = 5, 6$, then $\pi \neq (2^n)$.

Theorem 18. Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially C_5 -graphic if and only if the following conditions hold:

- (1) $d_5 \geq 2$. (2) For $i = 1, 2, d_1 = n - i$ implies $d_{4-i} \geq 3$
- (3) If $\pi = (d_1, d_2, 2^k, 1^{n-k-2})$, then $d_1 + d_2 \leq n + k - 2$.

Chen [2] characterized the potentially C_k -graphic sequences for each $k = 6$.

Theorem 19. Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 6$. Then π is potentially C_6 -graphic if and only if the following conditions hold:

- (1) $d_6 \geq 2$.
- (2) If $n = 7, 8$, then $\pi \neq (2^n)$.
- (3) For $i = 1, 2, 3$, $d_1 = n - i$ implies $d_{5-i} \geq 3$.
- (4) If $\pi = (d_1, d_2, 2^k, 1^{n-k-2})$, then $d_1 + d_2 \leq n + k - 2$; if $\pi = (d_1, d_2, 3, 2^k, 1^{n-k-3})$, then $d_1 + d_2 \leq n + k$; if $\pi = (d_1, d_2, 3, 3, 2^k, 1^{n-k-4})$, then $d_1 + d_2 \leq n + k + 2$.

Yin, Chen and Chen [60] characterized the potentially ${}_k C_l$ -graphic sequences for each $k = 3, 4 \leq l \leq 5$ and $k = 4, l = 5$.

Theorem 20. Let $\pi = (d_1, d_2, \dots, d_n) \in GS_n$ be a potentially C_4 -graphic sequence. Then π is potentially ${}_3 C_4$ -graphic if and only if π satisfies one of the following conditions:

- (1) $d_2 \geq 3$ and $\pi \neq (3^2, 2^4)$;
- (2) $\pi = (d_1, 2^k, 1^{n-k-1})$ with $2 \leq d_1 \leq 3$ and $k \geq 6$, and $\pi \neq (2^8)$ and (2^9) ;
- (3) $\pi = (d_1, 2^k, 1^{n-k-1})$ with $4 \leq d_1 \leq n - 2$ and $k \geq 5$, and $\pi \neq (4, 2^6)$ and $(4, 2^7)$.

Theorem 21. Let $\pi = (d_1, d_2, \dots, d_n) \in GS_n$ be a potentially C_5 -graphic sequence. Then π is potentially ${}_3 C_5$ -graphic if and only if π satisfies one of the following conditions:

- (1) $d_2 \geq 3$ and $\pi \neq (3^2, 2^4)$ and $(3^2, 2^5)$;
- (2) $\pi = (d_1, 2^k, 1^{n-k-1})$ with $2 \leq d_1 \leq 3$ and $k \geq 11$, and $\pi \neq (2^{13})$ and (2^{14}) ;
- (3) $\pi = (d_1, 2^k, 1^{n-k-1})$ with $4 \leq d_1 \leq 5$ and $k \geq 10$, and $\pi \neq (4, 2^{11})$ and $(4, 2^{12})$.
- (4) $\pi = (d_1, 2^k, 1^{n-k-1})$ with $6 \leq d_1 \leq n - 4$ and $k \geq 9$, and $\pi \neq (4, 2^{10})$ and $(4, 2^{11})$.

Theorem 22. Let $\pi = (d_1, d_2, \dots, d_n) \in GS_n$ be a potentially C_5 -graphic sequence. Then π is potentially ${}_4 C_5$ -graphic if and only if π satisfies one of the following conditions:

- (1) $d_2 \geq 3$;
- (2) $\pi = (d_1, 2^k, 1^{n-k-1})$ with $2 \leq d_1 \leq 3$ and $k \geq 8$, and $\pi \neq (2^{10})$ and (2^{11}) ;
- (3) $\pi = (d_1, 2^k, 1^{n-k-1})$ with $4 \leq d_1 \leq n - 4$ and $k \geq 7$, and $\pi \neq (4, 2^8)$ and $(4, 2^9)$.

Chen, Yin and Fan [10] characterized the potentially ${}_k C_l$ -graphic sequences for each $3 \leq k \leq 5, l = 6$.

Theorem 23. Let $\pi = (d_1, d_2, \dots, d_n) \in GS_n$, $n \geq 6$, and $\pi \neq (3^2, 2^{10}), (2^{19}), (2^{20}), (4, 2^{17}), (4, 2^{18}), (6, 2^{16}), (6, 2^{17}), (8, 2^{15}), (8, 2^{16})$. Then π is potentially ${}_3 C_6$ -graphic if and only if π be a potentially C_6 -graphic sequence, and π satisfies one of the following conditions:

- (1) $d_3 \geq 3$, and if $d_1 = d_3 = 3, d_4 = 2$, then $d_{10} = 2$;
- (2) $d_2 \geq 4, d_3 = 2, d_7 = 2$;

(3) $d_2 = 3, d_3 = 2$, and if $4 \geq d_1 \geq 3$, then $d_{10} = 2$, and if $n-4 \geq d_1 \geq 5$, then $d_9 = 2$.

(4) $d_2 = 2$, and if $3 \geq d_1 \geq 2$, then $d_{18} = 2$, and if $5 \geq d_1 \geq 4$, then $d_{17} = 2$, and if $7 \geq d_1 \geq 6$, then $d_{16} = 2$, and if $n-7 \geq d_1 \geq 8$, then $d_{15} = 2$.

Theorem 24. Let $\pi = (d_1, d_2, \dots, d_n) \in GS_n$, $n \geq 6$, and $\pi \neq (2^{16}), (2^{17}), (4, 2^{14}), (4, 2^{15}), (6, 2^{13}), (6, 2^{14})$. Then π is potentially ${}_4C_6$ -graphic if and only if π be a potentially C_6 -graphic sequence, and π satisfies one of the following conditions:

(1) $d_3 \geq 3$, and if $d_1 = d_3 = 3, d_4 = 2$, then $d_{10} = 2$;

(2) $d_2 \geq 4, d_3 = 2, d_7 = 2$;

(3) $d_2 = 3, d_3 = 2$, and if $4 \geq d_1 \geq 3$, then $d_{10} = 2$, and if $n-4 \geq d_1 \geq 5$, then $d_9 = 2$.

(4) $d_2 = 2$, and if $3 \geq d_1 \geq 2$, then $d_{15} = 2$, and if $5 \geq d_1 \geq 4$, then $d_{14} = 2$, and if $n-7 \geq d_1 \geq 6$, then $d_{13} = 2$.

Theorem 25. Let $\pi = (d_1, d_2, \dots, d_n) \in GS_n$, $n \geq 6$, and $\pi \neq (2^{12}), (2^{13}), (4, 2^{10}), (4, 2^{11})$. Then π is potentially ${}_5C_6$ -graphic if and only if π be a potentially C_6 -graphic sequence, and π satisfies one of the following conditions:

(1) $d_2 \geq 3$;

(2) $3 \geq d_1 \geq 2, d_2 = 2, d_{11} = 2$;

(3) $n-6 \geq d_1 \geq 4, d_2 = 2, d_{10} = 2$.

Luo and Warner [54] characterized the potentially K_4 -graphic sequences.

Theorem 26. Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence without zero terms and with $d_4 \geq 3$ and $n \geq 4$. Then π is potentially K_4 -graphic if and only if $d_4 \geq 3$ and $\pi \neq (n-1, 3^s, !^{n-s-1})$ for each $s = 4, 5$ except the following sequences:

$n = 5 : (4, 3^4), (3^4, 2)$;

$n = 6 : (4^6), (4^2, 3^4), (4, 3^4, 2), (3^6), (3^5, 1), (3^4, 2^2)$;

$n = 7 : (4^7), (4^3, 3^4), (4, 3^6), (4, 3^5, 1), (3^6, 2), 3^5, 2, 1)$;

$n = 8 : (3^7, 1), (3^6, 1^2)$.

Eschen and Niu [14] and Lai[31] independently characterized the potentially $K_4 - e$ -graphic sequences.

Theorem 27. Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 4$. Then π is potentially $(K_4 - e)$ -graphic if and only if the following conditions hold:

(1) $d_2 \geq 3$. (2) $d_4 \geq 2$

(3) If $n = 5, 6$, then $\pi \neq (3^2, 2^{n-2})$ and $\pi \neq (3^6)$.

Yin and Yin [73] characterized the potentially $K_5 - e$ and K_6 -graphic sequences.

Theorem 28 Let $n \geq 5$ and $\pi = (d_1, d_2, \dots, d_n) \in NS_n$ be a positive graphic sequence with $d_3 \geq 4$ and $d_5 \geq 3$. Then π is potentially

$K_5 - e$ -graphic if and only if π is not one of the following sequences:
 $(n-1, 4^6, 1^{n-7}), (n-1, 4^2, 3^4, 1^{n-7}), (n-1, 4^2, 3^3, 1^{n-6});$

$n = 6: (4^6), (4^4, 3^2), (4^3, 3^2, 2);$

$n = 7: (4^3, 3^4), (5^2, 4, 3^4), (4^7), (4^5, 3^2), (5, 4^3, 3^3), (5^2, 4^5), (5, 4^5, 3),$
 $(4^3, 3^2, 2^2), (4^4, 3^2, 2), (5, 4^2, 3^3, 2), (4^6, 2), (4^3, 3^3, 1);$

$n = 8: (5^8), (4^8), (5^2, 4^6), (6, 4^7), (4^4, 3^4), (5, 4^2, 3^5), (4^6, 3^2), (5, 4^6, 3),$
 $(4^3, 3^4, 2), (4^7, 2), (4^4, 3^3, 1), (5, 4^2, 3^4, 1), (4^3, 3^3, 2, 1), (4^6, 3, 1), (5, 4^6, 1);$

$n = 9: (4^9), (4^3, 3^5, 1), (4^8, 2), (4^7, 3, 1), (5, 4^7, 1), (4^3, 3^4, 1^2), (4^7, 1^2);$

$n = 10: (4^8, 1^2).$

Theorem 29 Let $n \geq 18$ and $\pi = (d_1, d_2, \dots, d_n) \in NS_n$ be a positive graphic sequence with $d_6 \geq 5$. Then π is potentially A_6 -graphic if and only if $\pi_6 \notin \{(2), (2^2), (3, 1), (3^2), (3, 2, 1), (3^2, 2), (3^3, 1), (3^2, 1^2)\}$.

Yin and Chen [61] characterized the potentially $K_{r,s}$ -graphic sequences for $r = 2, s = 3$ and $r = 2, s = 4$.

Theorem 30 Let $n \geq 5$ and $\pi = (d_1, d_2, \dots, d_n) \in GS_n$. Then π is potentially $K_{2,3}$ -graphic if and only if π satisfies the following conditions:

(1) $d_2 \geq 3$ and $d_5 \geq 2$;

(2) If $d_1 = n-1$ and $d_2 = 3$, then $d_5 = 3$;

(3) $\pi \neq (3^2, 2^4), (3^2, 2^5), (4^3, 2^3), (n-1, 3^5, 1^{n-6})$ and $(n-1, 3^6, 1^{n-7})$.

Theorem 31 Let $n \geq 6$ and $\pi = (d_1, d_2, \dots, d_n) \in GS_n$. Then π is potentially $K_{2,4}$ -graphic if and only if π satisfies the following conditions:

(1) $d_2 \geq 4$ and $d_6 \geq 2$;

(2) If $d_1 = n-1$ and $d_2 = 4$, then $d_3 = 4$ and $d_6 \geq 3$;

(3) $\pi \neq (4^3, 2^4), (4^2, 2^5), (4^2, 2^6), (5^2, 4, 2^4), (5^3, 3, 2^3), (6, 5^2, 2^5), (5^3, 2^4, 1),$
 $(6^3, 2^6), (n-1, 4^2, 3^4, 1^{n-7}), (n-1, 4^2, 3^5, 1^{n-8}), (n-2, 4^2, 2^3, 1^{n-6}),$ and
 $(n-2, 4^3, 2^2, 1^{n-6}).$

Chen [3] characterized the potentially $K_5 - 2K_2$ -graphic sequences for $5 \leq n \leq 8$. Hu and Lai [23] characterized the potentially $K_5 - P_3, K_5 - A_3, K_5 - K_3, K_5 - K_{1,3}$ and $K_5 - 2K_2$ -graphic sequences where A_3 is $P_2 \cup K_2$.

Theorem 32 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - P_3$ -graphic if and only if the following conditions hold:

(1) $d_1 \geq 4, d_3 \geq 3$ and $d_5 \geq 2$.

(2) $\pi \neq (4, 3^2, 2^3), (4, 3^2, 2^4)$ and $(4, 3^6)$.

Theorem 33 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - A_3$ -graphic if and only if the following conditions hold:

(1) $d_4 \geq 3$ and $d_5 \geq 2$.

(2) $\pi \neq (n-1, 3^3, 2^{n-k}, 1^{k-4})$ where $n \geq 6$ and $k = 4, 5, \dots, n-2, n$ and k have the same parity.

(3) $\pi \neq (3^4, 2^2), (3^6), (3^4, 2^3), (3^6, 2), (4, 3^6), (3^7, 1), (3^8), (n-1, 3^5, 1^{n-6})$ and $(n-1, 3^6, 1^{n-7})$.

Theorem 34 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - K_3$ -graphic if and only if the following conditions hold:

- (1) $d_2 \geq 4$ and $d_5 \geq 2$.
- (2) $\pi \neq (4^2, 2^4), (4^2, 2^5), (4^3, 2^3)$ and (4^6) .

Theorem 35 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - K_{1,3}$ -graphic if and only if the following conditions hold:

- (1) $d_1 \geq 4$ and $d_4 \geq 3$.
- (2) $\pi \neq (4, 3^4, 2), (4^6), (4^2, 3^4), (4, 3^6), (4^7), (4, 3^5, 1), (n-1, 3^4, 1^{n-5})$ and $(n-1, 3^5, 1^{n-6})$.

Theorem 36 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - 2K_2$ -graphic if and only if the following conditions hold:

- (1) $d_1 \geq 4$ and $d_5 \geq 3$;
- (2)

$$\pi \neq \begin{cases} (n-i, n-j, 3^{n-i-j-2k}, 2^{2k}, 1^{i+j-2}) \\ n-i-j \text{ is even;} \\ (n-i, n-j, 3^{n-i-j-2k-1}, 2^{2k+1}, 1^{i+j-2}) \\ n-i-j \text{ is odd.} \end{cases}$$

where $1 \leq j \leq n-5$ and $0 \leq k \leq \lfloor \frac{n-j-i-4}{2} \rfloor$.

- (3) $\pi \neq (4^2, 3^4), (4, 3^4, 2), (5, 4, 3^5), (5, 3^5, 2), (4^7), (4^3, 3^4), (4^2, 3^4, 2), (4, 3^6), (4, 3^5, 1), (4, 3^4, 2^2), (5, 3^7), (5, 3^6, 1), (4^8), (4^2, 3^6), (4^2, 3^5, 1), (4, 3^6, 2), (4, 3^5, 2, 1), (4, 3^7, 1), (4, 3^6, 1^2), (n-1, 3^5, 1^{n-6})$ and $(n-1, 3^6, 1^{n-7})$.

Hu and Lai [21] characterized the potentially $K_5 - C_4$ -graphic sequences.

Theorem 37 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $(K_5 - C_4)$ -graphic if and only if the following conditions hold:

- (1) $d_1 \geq 4$.
- (2) $d_5 \geq 2$.
- (3) $\pi \neq ((n-2)^2, 2^{n-2})$ for $n \geq 6$, where the symbol x^y stands for y consecutive terms x .
- (4) $\pi \neq (n-k, k+i, 2^i, 1^{n-i-2})$ where $i = 3, 4, \dots, n-2k$ and $k = 1, 2, \dots, \lfloor \frac{n-1}{2} \rfloor - 1$.
- (5) If $n = 6$, then $\pi \neq (4, 2^5)$.
- (6) If $n = 7$, then $\pi \neq (4, 2^6)$.

Hu and Lai [22] characterized the potentially $K_5 - Z_4$ -graphic sequences.

Theorem 38 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $(K_5 - Z_4)$ -graphic if and only if the following conditions hold:

- (1) $d_1 \geq 4, d_2 \geq 3$ and $d_4 \geq 2$.

Hu, Lai and Wang[25] characterized the potentially $K_5 - P_4$ and $K_5 - Y_4$ -graphic sequences where Y_4 is a tree on 5 vertices and 3 leaves.

Theorem 39 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - P_4$ -graphic if and only if the following conditions hold:

- (1) $d_2 \geq 3$.
- (2) $d_5 \geq 2$.
- (3) $\pi \neq (n-1, k, 2^t, 1^{n-2-t})$ where $n \geq 5$, $k, t = 3, 4, \dots, n-2$, and, k and t have different parities.
- (4) For $n \geq 5$, $\pi \neq (n-k, k+i, 2^i, 1^{n-i-2})$ where $i = 3, 4, \dots, n-2k$ and $k = 1, 2, \dots, \lfloor \frac{n-1}{2} \rfloor - 1$.
- (5) If $n = 6, 7$, then $\pi \neq (3^2, 2^{n-2})$.

Theorem 40 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_5 - Y_4$ -graphic if and only if the following conditions hold:

- (1) $d_3 \geq 3$.
- (2) $d_4 \geq 2$.
- (3) $\pi \neq (3^6)$.

Hu and Lai [24] characterized the potentially $K_{3,3}$ and $K_6 - C_6$ -graphic sequences.

Theorem 41 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 6$. Then π is potentially $K_{3,3}$ -graphic if and only if the following conditions hold:

- (1) $d_6 \geq 3$;
- (2) For $i = 1, 2$, $d_1 = n - i$ implies $d_{4-i} \geq 4$;
- (3) $d_2 = n - 1$ implies $d_3 \geq 5$ or $d_6 \geq 4$;
- (4) $d_1 + d_2 = 2n - i$ and $d_{n-i+3} = 1$ ($3 \leq i \leq n - 4$) implies $d_3 \geq 5$ or $d_6 \geq 4$;
- (5) $d_1 + d_2 = 2n - i$ and $d_{n-i+4} = 1$ ($4 \leq i \leq n - 3$) implies $d_3 \geq 4$;
- (6) $\pi = (d_1, d_2, 3^4, 2^t, 1^{n-6-t})$ or $(d_1, d_2, 4^2, 3^2, 2^t, 1^{n-6-t})$ implies $d_1 + d_2 \leq n + t + 2$;
- (7) $\pi = (d_1, d_2, 4, 3^4, 2^t, 1^{n-7-t})$ implies $d_1 + d_2 \leq n + t + 3$;
- (8) For $t = 5, 6$, $\pi \neq (n-i, k+i, 4^t, 2^{k-t}, 1^{n-2-k})$ where $i = 1, \dots, \lfloor \frac{n-k}{2} \rfloor$ and $k = t, \dots, n - 2i$;
- (9) $\pi \neq (5^4, 3^2, 2), (4^6), (3^6, 2), (6^4, 3^4), (4^2, 3^6), (4, 3^6, 2), (3^6, 2^2), (3^8), (3^7, 1), (4, 3^8), (4, 3^7, 1), (3^8, 2), (3^7, 2, 1), (3^9, 1), (3^8, 1^2), (n-1, 4^2, 3^4, 1^{n-7}), (n-1, 4^2, 3^5, 1^{n-8}), (n-1, 5^3, 3^3, 1^{n-7}), (n-2, 4, 3^5, 1^{n-7}), (n-2, 4, 3^6, 1^{n-8}), (n-3, 3^6, 1^{n-7}), (n-3, 3^7, 1^{n-8})$.

Theorem 42 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 6$. Then π is potentially $K_6 - C_6$ -graphic if and only if the following conditions hold:

- (1) $d_6 \geq 3$;
- (2) For $i = 1, 2$, $d_1 = n - i$ implies $d_{4-i} \geq 4$;

- (3) $d_2 = n - 1$ implies $d_4 \geq 4$;
- (4) $d_1 + d_2 = 2n - i$ and $d_{n-i+3} = 1$ ($3 \leq i \leq n - 4$) implies $d_4 \geq 4$;
- (5) $d_1 + d_2 = 2n - i$ and $d_{n-i+4} = 1$ ($4 \leq i \leq n - 3$) implies $d_3 \geq 4$;
- (6) $\pi = (d_1, d_2, d_3, 3^k, 2^t, 1^{n-3-k-t})$ implies $d_1 + d_2 + d_3 \leq n + 2k + t + 1$;
- (7) $\pi = (d_1, d_2, 3^4, 2^t, 1^{n-6-t})$ implies $d_1 + d_2 \leq n + t + 2$;
- (8) $\pi \neq (n - i, k, t, 3^t, 2^{k-i-t-1}, 1^{n-2-k+i})$ where $i = 1, \dots, \lfloor \frac{n-t-1}{2} \rfloor$ and $k = i + t + 1, \dots, n - i$ and $t = 4, 5, \dots, k - i - 1$;
- (9) $\pi \neq (3^6, 2), (4^2, 3^6), (4, 3^6, 2), (3^6, 2^2), (3^8), (3^7, 1), (4, 3^8), (4, 3^7, 1), (3^8, 2), (3^7, 2, 1), (3^9, 1), (3^8, 1^2), (n - 1, 4^2, 3^4, 1^{n-7}), (n - 1, 4^2, 3^5, 1^{n-8}), (n - 2, 4, 3^5, 1^{n-7}), (n - 2, 4, 3^6, 1^{n-8}), (n - 3, 3^6, 1^{n-7}), (n - 3, 3^7, 1^{n-8})$.

Xu and Hu[57] characterized the potentially $K_{1,4} + e$ -graphic sequences. Chen and Li [8] characterized the potentially $K_{1,t} + e$ -graphic sequences.

Theorem 43 Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $K_{1,4} + e$ -graphic if and only if $d_1 \geq 4, d_3 \geq 2$.

Theorem 44 Let $t \geq 3$, $\pi = (d_1, d_2, \dots, d_n)$ is a graphic sequence with $n \geq t + 1$. Then π is potentially $K_{1,t} + e$ -graphic if and only if $d_1 \geq t, d_3 \geq 2$.

Open Problems

Problem 1. Determine $\sigma(K_{r+1} - G, n)$ for G is a graph on k vertices and j edges which contains a graph $K_3 \cup K_{1,3}$ but not contains a cycle on 4 vertices and not contains Z_4, P_3 .

Problem 2. Determine $\sigma(K_{r+1} - G, n)$ for $G = K_3 \cup iK_2 \cup jP_2 \cup tK_3$.

Problem 3. Determine $\sigma(K_{r+1} - G, n)$ for graph G which contains C_3, C_4, \dots, C_l but not contains a cycle on $l + 1$ vertices ($4 \leq l \leq r$).

Problem 4. Determine $\sigma(K_{r+1} - G, n)$ for graph G which contains C_3, C_4, \dots, C_{r+1} .

Problem 5. Determine $\sigma(K_{r+1} - G, n)$ for small n .

Problem 6. Characterize potentially $K_{r+1} - G$ -graphic sequences for the remaining G .

Acknowledgment

The first author is particularly indebted to Professor Jiongsheng Li for introducing him to degree sequences. The authors wish to thank Professor Gang Chen, R.J. Gould, Jiongsheng Li, Rong Luo, John R. Schmitt, Zi-Xia Song, Amitabha Tripathi, Jianhua Yin and Mengxiao Yin for sending some their papers to us.

References

- [1] B. Bollobás, Extremal Graph Theory, Academic Press, London, 1978.
- [2] Chen Gang, Potentially C_6 -graphic sequences, J. Guangxi Univ. Nat. Sci. Ed. 28 (2003), no. 2, 119–124.
- [3] Chen Gang, Characterize the potentially $K_{1,2,2}$ -graphic sequences, Journal of Qingdao University of Science and Technology, 27(2006), 86–88.
- [4] Chen Gang, The smallest degree sum that yields potentially fan graphical sequences, Xibei Shifan Daxue Xuebao Ziran Kexue Ban 42 (2006), no. 4, 27–30.
- [5] Chen Gang, An extremal problem on potentially $K_{r,s,t}$ -graphic sequences, J. YanTai University, 19(4) 2006,245-252.
- [6] Chen Gang, Potentially $K_{3,s}-ke$ graphical sequences, Guangxi Sciences 13(3)(2006), 164-171.
- [7] Chen Gang, A note on potentially $K_{1,1,t}$ -graphic sequences, Australasian Journal of Combinatorics, 37(2007), 21-26.
- [8] Chen Gang, Li Xining, On potentially $K_{1,t}+e$ -graphic sequences, J. Zhangzhou Teach. Coll., 20 (2007), no. 3, 5–7.
- [9] Chen Gang, Yin Jianhua, The smallest degree sum that yields potentially W_5 -graphic sequences, J. XuZhou Normal University, 21(2) 2003,5-7.
- [10] Chen Gang, Yin Jianhua, Fan Yingmei, Potentially kC_6 -graphic sequences. J. Guangxi Norm. Univ. Nat. Sci. 24 (2006), no. 3, 26–29.
- [11] Guantao Chen, Michael Ferrara, Ronald J.Gould, John R. Schmitt, Graphic sequences with a realization containing a complete multipartite subgraph, accepted by Discrete Mathematics.
- [12] P.Erdős, T.Gallai, Graphs with given degrees of vertices, Math. Lapok,11(1960),264-274.
- [13] P.Erdős, M.S.Jacobson and J. Lehel, Graphs realizing the same degree sequences and their respective clique numbers, in Graph Theory, Combinatorics and Application, Vol. 1(Y. Alavi et al., eds.), John Wiley and Sons, Inc., New York, 1991, 439-449.
- [14] Elaine M.Eschen, Jianbing Niu, On potentially K_4-e -graphic sequences, Australasian Journal of Combinatorics, 29(2004), 59-65.

- [15] Michael J.Ferrara, Graphic sequences with a realization containing a union of cliques, *Graphs and Combinatorics* 23 (2007) 263-269.
- [16] M.Ferrara, R.Gould, J.Schmitt, Potentially K_s^t -graphic degree sequences, submitted.
- [17] M.Ferrara, R.Gould, J.Schmitt, Graphic sequences with a realization containing a friendship graph, *Ars Combinatoria* 85 (2007), 161-171.
- [18] M.Ferrara, M.Jacobson, J.Schmitt, M.Siggers, Potentially H -bigraphic sequences, submitted.
- [19] Ronald J.Gould, Michael S.Jacobson, J.Lehel, Potentially G -graphical degree sequences. *Combinatorics, graph theory, and algorithms*, Vol. I, II (Kalamazoo, MI, 1996), 451-460, New Issues Press, Kalamazoo, MI, 1999.
- [20] Gautam Gupta, Puneet Joshi, Amitabha Tripathi, Graphic sequences of trees and a problem of Frobenius, *Czechoslovak Mathematical Journal*, 57(2007): 49-52.
- [21] Hu Lili, Lai Chunhui, On potentially $K_5 - C_4$ -graphic sequences, accepted by *Ars Combinatoria*.
- [22] Hu Lili, Lai Chunhui, On potentially $K_5 - Z_4$ -graphic sequences, submitted.
- [23] Hu Lili, Lai Chunhui, On potentially $K_5 - E_3$ -graphic sequences, accepted by *Ars Combinatoria*.
- [24] Hu Lili, Lai Chunhui, On Potentially 3-regular graph graphic Sequences, accepted by *Utilitas Mathematica*.
- [25] Hu Lili, Lai Chunhui, Wang Ping , On potentially $K_5 - H$ -graphic sequences, accepted by *Czechoslovak Mathematical Journal*.
- [26] Huang Qin, On potentially $K_m - e$ -graphic sequences, *J. ZhangZhou Teachers College*, 15(4) 2002, 26-28.
- [27] Huang Qin, On potentially $K_5 - e$ -graphic sequences, *J. XinJiang University*, 22(3) 2005, 276-284.
- [28] D.J. Kleitman, D.L. Wang , Algorithm for constructing graphs and digraphs with given valences and factors,*Discrete Math.*, 6(1973),79-88.
- [29] Lai Chunhui, On potentially C_k -graphic sequences, *J. ZhangZhou Teachers College*, 11(4) 1997, 27-31.

- [30] Lai Chunhui, A note on potentially $K_4 - e$ graphical sequences, Australasian J. of Combinatorics 24,2001, 123-127.
- [31] Lai Chunhui, Characterize the potentially $K_4 - e$ graphical sequences, J. ZhangZhou Teachers College, 15(3) 2002, 53-59.
- [32] Lai Chunhui, The Smallest Degree Sum that Yields Potentially C_k -graphic Sequences, J.Combin. Math. Combin. Comput. 49(2004), 57-64.
- [33] Lai Chunhui, An extremal problem on potentially $K_{p,1,\dots,1}$ -graphic sequences, J.Zhangzhou Teachers College 17(4) (2004),11-13.
- [34] Lai Chunhui, An extremal problem on potentially $K_{p,1,1}$ -graphic sequences, Discrete Mathematics and Theoretical Computer Science 7(2005), 75-80.
- [35] Lai Chunhui, An extremal problem on potentially $K_m - C_4$ -graphic sequences, Journal of Combinatorial Mathematics and Combinatorial Computing, 61 (2007), 59-63.
- [36] Lai Chunhui, An extremal problem on potentially $K_m - P_k$ -graphic sequences, accepted by International Journal of Pure and Applied Mathematics.
- [37] Lai Chunhui, The smallest degree sum that yields potentially $K_{r+1} - Z$ -graphical Sequences, accepted by Ars Combinatoria.
- [38] Lai Chunhui, Hu Lili, An extremal problem on potentially $K_{r+1} - H$ -graphic sequences, accepted by Ars Combinatoria.
- [39] Lai Chunhui, Sun Yuzhen, An extremal problem on potentially $K_{r+1} - (kP_2 \cup tK_2)$ -graphic sequences, submitted.
- [40] Lai Chunhui, Yan Guiying, On potentially $K_{r+1} - U$ -graphical Sequences, accepted by Utilitas Mathematica.
- [41] Li Jiongsheng, Luo Rong, Potentially ${}_3C_l$ -Graphic Sequences, J. Univ. Sci. Technol. China, 29(1) 1999: 1-8.
- [42] Li Jiongsheng, Luo Rong, Liu Yunkai, An extremal problem on potentially ${}_3C_l$ -graphic sequences, J.Math.Study, 31(4)(1998), 362-369.
- [43] Li Jiongsheng, Song Zi-xia, The smallest degree sum that yields potentially P_k -graphical sequences, J. Graph Theory,29(1998), 63-72.
- [44] Li Jiongsheng, Song Zi-xia, On the potentially P_k -graphic sequences, Discrete Math. 195(1999), 255-262.

- [45] Li Jiongsheng, Song Zi-xia, An extremal problem on the potentially P_k -graphic sequences, *Discrete Math.*, 212(2000), 223-231.
- [46] Li Jiongsheng, Song Zi-xia, Luo Rong, The Erdős-Jacobson-Lehel conjecture on potentially P_k -graphic sequence is true, *Science in China(Series A)*, 41(5)(1998), 510-520.
- [47] Li Jiongsheng, Song Zi-xia, Wang Ping, The Erdos-Jacobson-Lehel conjecture about potentially P_k -graphic sequences, *J. China Univ. Sci. Tech.* 28 (1998), no. 1, 1-9.
- [48] Li Jiongsheng, Yin Jianhua, A variation of an extremal theorem due to Woodall, *Southeast Asian Bull. Math.* 25(2001), 427-434.
- [49] Li Jiongsheng, Yin Jianhua, On potentially $A_{r,s}$ -graphic sequences. *J. Math. Study* 34 (2001), no. 1, 1-4.
- [50] Li Jiongsheng, Yin Jianhua, Extremal graph theory and degree sequences, *Adv. Math.* 33(2004), 273-283.
- [51] Li Jiong Sheng, Yin Jianhua, The threshold for the Erdos, Jacobson and Lehel conjecture to be true. *Acta Math. Sin. (Engl. Ser.)* 22 (2006), no. 4, 1133-1138.
- [52] Liu Mingjing, Hu Lili, On Potentially $K_5 - Z_5$ graphic sequences, *J. Zhangzhou Normal University*, 57(3)(2007), 20-24.
- [53] Luo Rong, On potentially C_k -graphic sequences, *Ars Combinatoria* 64(2002), 301-318.
- [54] Luo Rong, Warner Morgan , On potentially K_k -graphic sequences, *Ars Combin.* 75(2005), 233-239.
- [55] Mubayi Dhruv. Graphic sequences that have a realization with large clique number. *J. Graph Theory* 34 (2000), no. 1, 20-29.
- [56] J.R.Schmitt, M.Ferrara, An Erdős-Stone Type Conjecture for Graphic Sequences, *Electronic Notes in Discrete Mathematics*, 28(2007), 131-135.
- [57] Xu Zhenghua, Hu Lili, Characterize the potentially $K_{1,4} + e$ graphical sequences, *ZhangZhou Teachers College*, 55(1) 2007, 4-8.
- [58] Yin Jianhua, The smallest degree sum that yields potentially $K_{1,1,3}$ -graphic sequences, *J. HaiNan University*, 22(3) 2004, 200-204.
- [59] Yin Jianhua, Some new conditions for a graphic sequence to have a realization with prescribed clique size, submitted.

- [60] Yin Jianhua, Chen Gang, Guoliang Chen, On potentially $_kC_l$ -graphic sequences. J. Combin. Math. Combin. Comput. 61 (2007), 141–148.
- [61] Yin Jianhua, Chen Gang, On potentially K_{r_1, r_2, \dots, r_m} -graphic sequences, Util. Math. 72 (2007), 149–161.
- [62] Yin Jianhua, Chen Gang, Schmitt John R. , Graphic Sequences with a realization containing a generalized Friendship Graph, Discrete Mathematics (2008), doi:10.1016/j.disc.2007.11.075, in press.
- [63] Yin Jianhua, Li Jiongsheng, The smallest degree sum that yields potentially $K_{r,r}$ -graphic sequences, Science in China Ser A, 2002, 45:694-705.
- [64] Yin Jianhua, Li Jiongsheng, On the threshold in the Erdos-Jacobson-Lehel problem, Math. Appl., 15(1)(2002), 123-128.
- [65] Yin Jianhua, Li Jiongsheng, An extremal problem on the potentially $K_{r,s}$ -graphic sequences, Discrete Math, 2003, 260:295-305.
- [66] Yin Jianhua, Li Jiongsheng, Two sufficient conditions for a graphic sequence to have a realization with prescribed clique size, Discrete Math., 301(2005) 218-227.
- [67] Yin Jianhua, Li Jiongsheng, Potentially $K_{r_1, r_2, \dots, r_l, r, s}$ -graphic sequences, Discrete Mathematics, 307(2007) 1167-1177.
- [68] Yin Jianhua, Li Jiongsheng, Chen Guoliang, The smallest degree sum that yields potentially $_3C_l$ -graphic sequences, Discrete Mathematics, 270(2003), 319-327.
- [69] Yin Jianhua, Li Jiongsheng, Chen Guoliang, A variation of a classical Turn-type extremal problem, Eur. J. Comb. 25(2004), 989-1002.
- [70] Jianhua Yin, Jiongsheng Li and Guoliang Chen, The smallest degree sum that yields potentially $K_{2,s}$ -graphic sequences, Ars Combinatoria, 74(2005), 213-222.
- [71] Yin Jianhua, Li Jiongsheng, Mao Rui, An extremal problem on the potentially $K_{r+1} - e$ -graphic sequences, Ars Combinatoria, 74(2005), 151-159.
- [72] Yin Mengxiao, The smallest degree sum that yields potentially $K_{r+1} - K_3$ -graphic sequences, Acta Math. Appl. Sin. Engl. Ser. 22(2006), no. 3, 451-456.
- [73] Yin Mengxiao, Yin Jianhua, On potentially H -graphic sequences, Czechoslovak Mathematical Journal, 57(2)(2007), 705-724.